

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
31 October 2002 (31.10.2002)

PCT

(10) International Publication Number
WO 02/086078 A2

(51) International Patent Classification⁷: **C12N**
(21) International Application Number: **PCT/US02/12655**
(22) International Filing Date: **22 April 2002 (22.04.2002)**
(25) Filing Language: **English**
(26) Publication Language: **English**
(30) Priority Data:
60/285,745 23 April 2001 (23.04.2001) **US**
60/338,955 10 December 2001 (10.12.2001) **US**

(71) Applicant (for all designated States except US): **UNIVERSITY OF VIRGINIA PATENT FOUNDATION**
[US/US]; Suite 1-110, 1224 West Main Street, Charlottesville, VA 22903 (US).

(72) Inventor; and

(75) Inventor/Applicant (for US only): **BROWN, Milton, L.** [US/US]; 614 Nettle Court, Charlottesville, VA 22903 (US).

(74) Agent: **BREEN, John, P.**; University of Virginia Patent Foundation, Suite 1-110, 1224 West Main Street, Charlottesville, VA 22903 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.


(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report


For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: **SYNTHESIS AND EVALUATION OF NOVEL PHTHALIMIDE MIMICS AS ANTI-ANGIOGENIC AGENTS**



(I)

(57) Abstract: The present invention is directed to novel thalidomide derivative compounds that have activity as anti-angiogenic compounds. More particularly the compounds have the general structure: where R₁ is selected from the group consisting of H, halo, alkyl, haloalkyl, -NH₂, hydroxy and alkoxy; R₂ is selected from the group consisting of optionally substituted bicyclic, optionally substituted aryl, and R₆ is H, or C₁-C₈ alkyl; R₁₉ is optionally substituted aryl; and m is 0-6.



(II)

WO 02/086078 A2

**Synthesis and Evaluation of Novel Phthalimide Mimics
as Anti-angiogenic Agents**

Related Applications

5 This application claims priority under 35 USC §119(e) to US
Provisional Application Serial Nos. 60/285,745, filed April 23, 2001 and 60/338,955,
filed December 10, 2001, the disclosures of which are incorporated herein.

Field of the Invention

10 The present invention is directed to novel substituted phthalimide and
isoquinoline derivatives, and the use of such derivatives as therapeutic agents. These
compounds have been discovered to have anti-angiogenic activity.

Background of the Invention

15 Inappropriate angiogenesis is the cause or an aggravating factor in
numerous disease states. For example, age related macular degeneration (ARMD)
refers to a condition that steals away central vision but leaves peripheral (side) vision
intact. This disease can be present in several forms and affects approximately 1 out of
5 individuals over the age of 65 and 1 out of 4 over the age of 75. That represents
20 approximately 20 million Americans.

 There are two forms of macular degeneration: dry macular
degeneration and wet macular degeneration. The dry form, in which the cells of the
macula slowly begin to break down, is diagnosed in 90 percent of macular
degeneration cases. It may occur in one eye or both. The wet form, although it only
25 accounts for 10 percent of the cases, results in 90 percent of the blindness. As the wet
form worsens, patients begin to have abnormal blood vessels growing behind the
macula. These vessels are very fragile and will leak fluid and blood (hence 'wet'
macular degeneration), causing rapid damage to the macula. As of yet no specific
pharmacological treatments are available.

30 The wet form of ARMD and other ocular diseases caused by the
formation of new blood vessels (neovascularization) are among the leading causes of
blindness. Thus there is a great need for the discovery of new drugs to treat such

-2-

ocular diseases. One approach is to administer pharmaceutical anti-angiogenic agents to prevent the inappropriate neovascularization.

Novel anti-angiogenic compounds also have utility as anti-cancer agents. Malignancies are characterized by the growth and spread of tumors. One
5 important factor in the progression of this disease is angiogenesis, a complex process in which capillary blood vessels grow in an ordered sequence of events. Once a tumor has started, every increase in tumor cell population must be preceded by an increase in new capillaries that converge on the tumor and supply the cells with oxygen and nutrients. Tumors may thus remain harmless and confined to their tissue of origin, as
10 long as an accompanying angiogenic program is prevented from being activated. Since the angiogenesis-dependent step in tumor progression is shared by solid tumors of all etiologies, the ability to inhibit tumor-associated angiogenesis is a promising approach in combating cancer.

A substantial body of experimental evidence supports the hypothesis
15 that tumor angiogenesis is fundamental for the growth and metastasis of solid tumors [Folkman, J. Natl. Cancer Inst., Vol. 82, pp. 4-6 (1989); N. Weidner, et al., Amer. J. Pathol., Vol. 143, pp. 401-409 (1993)]. Indeed, the majority of solid tumors are not even clinically detectable until after the occurrence of neovascularization, whose induction in solid tumors is mediated by one or more angiogenic factors.

20 Furthermore, angiogenesis is also important in a number of other pathological processes, including arthritis, psoriasis, diabetic retinopathy, chronic inflammation, scleroderma, hemangioma, retrolental fibroplasia and abnormal capillary proliferation in hemophiliac joints, prolonged menstruation and bleeding, and other disorders of the female reproductive system.

25 Accordingly, there is a need for compounds that have activity as anti-angiogenic agents and can be safely administered to patients to treat angiogenic-associated diseases. The present invention relates to a composition, comprising an anti-angiogenic compound, for use in treating angiogenic-associated diseases, as well as malignancies, including inhibition of primary tumor growth, tumor progression and
30 metastasis. More particularly, the present invention is directed to thalidomide derivatives and their use as anti-angiogenic compositions.

Thalidomide was originally prescribed as a sedative, however its use was discontinued when it was found to be a potent teratogen, causing serious birth defects, especially affecting limb development. The dysmelia (limb defects) seen with thalidomide is proposed to be caused by an inhibition of blood vessel growth in the developing fetal limb bud. Although this affect is vasculogenic (affecting the formation of a capillary bed), studies have demonstrated that thalidomide is also anti-angiogenic (affecting the formation of new blood vessels from sprouts of pre-existing vessels). Thalidomide is relatively nontoxic, when taken by nonpregnant adults and is now in phase 2 clinical trials as a potential anti-cancer agent as well as a treatment for vascular eye diseases such as diabetic retinopathy, retinopathy of prematurity, and macular degeneration.

Thalidomide has been reported by Folkman, et. al. (PNAS, 91(9):4082-5, 1994) as having significant anti-angiogenic efficacy. The effects of thalidomide on corneal angiogenesis induced by vascular endothelial growth factor (VEGF), has been reported (Kruse et al, Graefes Archive for Clinical & Experimental Ophthalmology, 236(6):461-6, 1998). In Kruse et al, corneal neovascularization was induced in rabbits by an intrastromal pellet loaded with 500 or 750 ng VEGF. Animals receiving two daily feedings of 200 mg/kg of thalidomide, responded with statistically significant inhibition ($P < 0.0001$) of corneal angiogenesis after the 5th day of treatment. This observation indicates that thalidomide has a significant anti-angiogenic effect against VEGF-induced ocular neovascular growth.


Thalidomide has also demonstrated inhibitory effects on angiogenesis in the basic fibroblast growth factor (bFGF) induced rabbit corneal micropocket assay and orally in mice models (Joussen et al. Graefes Archive for Clinical & Experimental Ophthalmology, 237(12):952-61, 1999 and Kenyon et al., Experimental Eye Research, 64(6):971-8, 1997). Thalidomide and a thalidomide analog (cc-1069) have been reported to inhibit the *in vitro* proliferation of endothelial cells (cells which make up the vascular system). The results of this study revealed a significant decrease in endothelial cell proliferation in cultures treated with thalidomide and/or cc-1069. Taken together, these data support a strong correlation between the anti-angiogenic potential and inhibition of endothelial cell proliferation for thalidomide.

-4-

Studies reveal that the S(-)-enantiomer of thalidomide has the strongest anti-angiogenic activity in both VEGF-induced and bFGF-induced corneal neovascularization. This enantioselective preference lends support to a possible receptor mediated mechanism. The present invention is directed to a novel series of thalidomide analogs and the use of such analogs as inhibitors of angiogenesis. More particularly, the thalidomide analogs of the present invention lack the piperidine-2,6-dione moiety of thalidomide.


Summary of the Invention

The present invention is directed to compounds having a general structure:



wherein R_1 is selected from the group consisting of H, halo, alkyl, haloalkyl, $-NR_5R_6$, hydroxy and alkoxy;

R_2 is selected from the group consisting of optionally substituted bicyclic, optionally substituted aryl, and



R_3 is selected from the group consisting of alkyl, phenyl, substituted phenyl, benzyl and substituted benzyl;

R_8 is selected from the group consisting of H or C_1 - C_6 alkyl, or R_8 and R_3 taken together, can form, with the adjacent ring, an optionally substituted 5- or 6-membered aryl;

R_5 and R_6 are independently H, or C_1 - C_8 alkyl;

R₁₉ is optionally substituted aryl; and m is 0-6. The present invention also encompasses compositions comprising those compounds and the use of such compositions for inhibiting angiogenesis.

5 Detailed Description of the Invention

Definitions

In describing and claiming the invention, the following terminology will be used in accordance with the definitions set forth below.

As used herein, the term "purified" and like terms relate to the isolation
10 of a molecule or compound in a form that is substantially free of contaminants normally associated with the molecule or compound in a native or natural environment.

As used herein, the term "treating" includes prophylaxis of the specific disorder or condition, or alleviation of the symptoms associated with a specific
15 disorder or condition and/or preventing or eliminating said symptoms.

As used herein, the term "halogen" or "halo" means Cl, Br, F, and I. Especially preferred halogens include Cl, Br, and F. The term "haloalkyl" as used herein refers to a C₁-C₄ alkyl radical bearing at least one halogen substituent, for example, chloromethyl, fluoroethyl or trifluoromethyl and the like.

20 The term "C₁-C_n alkyl" wherein n is an integer, as used herein, represents a branched or linear alkyl group having from one to the specified number of carbon atoms. Typically C₁-C₆ alkyl groups include, but are not limited to, methyl, ethyl, n-propyl, iso-propyl, butyl, iso-butyl, sec-butyl, tert-butyl, pentyl, hexyl and the like.

25 The term "C₂-C_n alkenyl" wherein n is an integer, as used herein, represents an olefinically unsaturated branched or linear group having from 2 to the specified number of carbon atoms and at least one double bond. Examples of such groups include, but are not limited to, 1-propenyl, 2-propenyl, 1,3-butadienyl, 1-butenyl, hexenyl, pentenyl, and the like.

30 The term "C₂-C_n alkynyl" wherein n is an integer refers to an unsaturated branched or linear group having from 2 to the specified number of carbon

atoms and at least one triple bond. Examples of such groups include, but are not limited to, 1-propynyl, 2-propynyl, 1-butylnyl, 2-butylnyl, 1-pentylnyl, and the like.

The term " C_3 - C_n cycloalkyl" wherein n is an integer refers to cyclic non-aryl group, for example C_3 - C_8 cycloalkyl, represents cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, and cyclooctyl.

The term "lower alkyl" as used herein refers to branched or straight chain alkyl groups comprising one to eight carbon atoms, including methyl, ethyl, propyl, isopropyl, n-butyl, t-butyl, neopentyl and the like.

As used herein, the term "optionally substituted" refers to from zero to four substituents, wherein the substituents are each independently selected. Each of the independently selected substituents may be the same or different than other substituents.

As used herein the term "aryl" refers to a mono- or bicyclic carbocyclic ring system having one or two aromatic rings including, but not limited to, phenyl, benzyl, naphthyl, tetrahydronaphthyl, indanyl, indenyl, and the like. "Optionally substituted aryl" includes aryl compounds having from zero to four substituents, and "substituted aryl" includes aryl compounds having one to three substituents, wherein the substituents, including alkyl, halo or amino substituents. The term (C_3 - C_8 alkyl)aryl refers to any aryl group which is attached to the parent moiety via the alkyl group.

The term "heterocyclic group" refers to a mono- or bicyclic carbocyclic ring system containing from one to three heteroatoms wherein the heteroatoms are selected from the group consisting of oxygen, sulfur, and nitrogen.

As used herein the term "heteroaryl" refers to a mono- or bicyclic carbocyclic ring system having one or two aromatic rings containing from one to three heteroatoms and includes, but is not limited to, furyl, thienyl, pyridyl and the like.

The term "bicyclic" represents either an unsaturated or saturated stable 7- to 12-membered bridged or fused bicyclic carbon ring. The bicyclic ring may be attached at any carbon atom which affords a stable structure. The term includes, but is not limited to, naphthyl, dicyclohexyl, dicyclohexenyl, and the like.

As used herein, the term "pharmaceutically acceptable carrier" encompasses any of the standard pharmaceutical carriers, such as a phosphate

buffered saline solution, water and emulsions such as an oil/water or water/oil emulsion, and various types of wetting agents.

As used herein, "effective amount" means an amount sufficient to produce a selected effect. For example, an effective amount of the phthalimide derivative is an amount of the compound sufficient to decrease endothelial cell proliferation, or decrease the growth rate of blood vessels, either *in vivo* or *in vitro*.

The term, "parenteral" means not through the alimentary canal but by some other route such as subcutaneous, intramuscular, intraspinal, or intravenous.

As used herein, the term "angiogenesis" means the generation of new blood vessels into a tissue or organ.


As used herein, the term "angiogenic-associated" disease or condition refers to a disease state or condition that is caused by, or aggravated by, inappropriate or excessive angiogenesis. For example, diseases that are considered angiogenic-associated include cancer as well as vascular eye diseases such as diabetic retinopathy, retinopathy of prematurity, and macular degeneration.

Compounds of the present invention that have one or more asymmetric carbon atoms may exist as the optically pure enantiomers, or optically pure diastereomers, as well as mixtures of enantiomers, mixtures of diastereomers, and racemic mixtures of such stereoisomers. The present invention includes within its scope all such isomers and mixtures thereof.


The Invention

The present invention relates a novel series of substituted phthalimide and isoquinoline derivatives that are anticipated to have anti-angiogenic activity. More particularly, the invention is directed to a series of thalidomide derivatives wherein the piperidine-2,6-dione moiety has been replaced with other structures as shown below:

-8-




In accordance with one embodiment a novel compound is provided having a general structure selected from the group consisting of:




wherein R_1 is selected from the group consisting of H, halo, C_1 - C_8 alkyl, C_1 - C_8 haloalkyl, $-NR_5R_7$, hydroxy and C_1 - C_8 alkoxy;

R_2 is selected from the group consisting of optionally substituted bicyclic, optionally substituted aryl, and

-9-



5 R_3 is selected from the group consisting of



10 R_8 is selected from the group consisting of H, C_1 - C_8 alkyl, or R_3 and R_8 taken together, can form, with the adjacent ring, an optionally substituted 5- or 6-membered aromatic ring;


R_{10} and R_{11} are independently selected from the group consisting of H, halo, C_1 - C_8 alkyl, C_1 - C_8 haloalkyl, $-\text{NR}_3\text{R}_7$, hydroxy and C_1 - C_8 alkoxy or R_{10} and R_{11} taken
15 together, can form, with the adjacent ring, an optionally substituted 5- or 6-membered aromatic ring;

R_5 and R_7 are independently H, or C_1 - C_8 alkyl;

R_6 is selected from the group consisting of H, or C_1 - C_{12} alkyl, C_2 - C_8 alkenyl
and C_2 - C_8 alkynyl;

20 n is an integer ranging from 0-2; and m is an integer ranging from 0-6.


In accordance with one embodiment the anti-angiogenic compound has the general structure:




-10-

wherein R_1 is selected from the group consisting of H, halo, C_1 - C_4 alkyl, carboxy and C_1 - C_8 alkoxy;

R_2 is selected from the group consisting of



R_3 is selected from the group consisting of



R_8 is selected from the group consisting of H, C_1 - C_8 alkyl, or R_3 and R_8 taken together, can form, with the adjacent ring, an optionally substituted 5- or 6-membered heteroaryl;

20 R_4 is selected from the group consisting of H, C_1 - C_8 alkyl, phenyl and benzyl;

R_{13} and R_9 are independently selected from the group consisting of H, halo and C_1 - C_8 alkoxy or R_{13} and R_9 taken together, can form with the adjacent ring, an optionally substituted C_5 - C_6 cycloalkyl or optionally substituted C_5 - C_6 aromatic ring;

25 R_{10} and R_{11} are independently selected from the group consisting of H, halo, C_1 - C_8 alkyl, C_1 - C_8 haloalkyl, $-NR_5R_7$, hydroxy and C_1 - C_8 alkoxy or R_{10} and R_{11} taken together, can form with the adjacent ring, an optionally substituted C_5 - C_6 cycloalkyl or an optionally substituted C_5 - C_6 aromatic ring;

R_5 and R_7 are independently H, or C_1 - C_8 alkyl;


30 R_6 is selected from the group consisting of H, or C_1 - C_{12} alkyl, C_2 - C_8 alkenyl and C_2 - C_8 alkynyl;

R_{12} is selected from the group consisting of H, halo, C_1 - C_4 haloalkyl, $-NR_5R_6$, C_1 - C_8 alkyl, hydroxy and C_1 - C_8 alkoxy;


-11-

n is an integer from 1-3 and m is an integer ranging from 0-6. In one embodiment, R_1 , R_4 and R_8 are each H, n is 1 and m is 0.


In accordance with one embodiment the anti-angiogenic compound has the general structure:



wherein R_1 is selected from the group consisting of H, halo, C_1 - C_8 alkyl, hydroxy and C_1 - C_8 alkoxy and R_3 is selected from the group consisting of




wherein R_6 is H, or C_1 - C_8 alkyl; R_{10} and R_{11} are independently selected from the group consisting of H, halo, C_1 - C_8 alkyl, C_1 - C_8 haloalkyl, $-NR_5R_7$, hydroxy and C_1 - C_8 alkoxy, or R_{10} and R_{11} taken together, can form with the adjacent ring, an optionally substituted C_5 - C_6 cycloalkyl or C_5 - C_6 aromatic ring; R_5 and R_7 are independently H, or C_1 - C_8 alkyl; and m is an integer ranging from 0-6. In one embodiment, R_1 is H and R_3 is selected from the group consisting of phenyl;



O=C1C(=O)N(R2)c2ccccc21R1

R_2 is selected from the group consisting of



20 wherein R₁₀ and R₁₁ are independently selected from the group consisting of H, halo, C₁-C₈ alkyl, C₁-C₈ haloalkyl, -NR₅R₇, hydroxy and C₁-C₈ alkoxy or R₁₀ and R₁₁ taken together, can form with the adjacent ring, an optionally substituted C₅-C₆ cycloalkyl or optionally substituted C₅-C₆ aromatic ring;

m is an integer ranging from 0-3;

25 R₅ and R₇ are independently H, or C₁-C₈ alkyl;


R₆ is selected from the group consisting of H, C₁-C₁₂ alkyl, C₂-C₈ alkenyl and C₂-C₈ alkynyl;

R_{12} is selected from the group consisting of H, halo, C_1 - C_4 haloalkyl, $-NR_5R_6$, C_1 - C_8 alkyl, hydroxy and C_1 - C_8 alkoxy; and


30 n is an integer ranging from 1-3.

-13-

In one embodiment the compound of the present invention has the general structure:



wherein R_2 is selected from the group consisting of




wherein R_{10} and R_{11} are independently selected from the group consisting of H, halo, C_1 - C_4 haloalkyl, $-NR_5R_7$, hydroxy and C_1 - C_8 alkoxy; and

R_5 and R_7 are independently H, or C_1 - C_4 alkyl;

R_{12} is selected from the group consisting of H, halo and C_1 - C_8 alkoxy; and


R_6 is C_1 - C_8 alkyl. In one preferred embodiment R_2 is selected from the group consisting of



wherein R_{10} and R_{11} are independently selected from the group consisting of H, halo, and C_1 - C_8 alkoxy.

-14-

In one preferred embodiment the compound has the structure:




5


wherein R_6 is C_1 - C_8 alkyl and R_{10} and R_{11} are independently selected from the group consisting of H and C_1 - C_4 alkoxy, or R_{10} and R_{11} taken together, can form, with the adjacent ring, an optionally substituted 5- or 6-membered aromatic ring.

One aspect of the present invention is directed to a method of treating a
 10 angiogenic-associated disease or condition. More particularly, one embodiment of the present invention is directed to inhibiting undesired angiogenesis in a warm-blooded vertebrate, including humans. The method comprises the step of administering to the human or animal a composition comprising an effective amount of a compound of the general structure:

15



20




25

wherein R_1 is selected from the group consisting of H, halo, C_1 - C_8 alkyl, C_1 - C_8 haloalkyl, $-NR_5R_7$, hydroxy and C_1 - C_8 alkoxy;


R_2 is selected from the group consisting of optionally substituted bicyclic, optionally substituted aryl, and

30



-15-

R_3 is selected from the group consisting of



5

R_8 is selected from the group consisting of H and $\text{C}_1\text{--C}_8$ alkyl, or R_3 and R_8 taken together, can form with the adjacent ring, an optionally substituted 5- or 6-membered aromatic ring;

R_{10} and R_{11} are independently selected from the group consisting of H, halo, $\text{C}_1\text{--C}_8$ alkyl, $\text{C}_1\text{--C}_8$ haloalkyl, $\text{---NR}_5\text{R}_7$, hydroxy and $\text{C}_1\text{--C}_8$ alkoxy or R_{10} and R_{11} taken together, can form, with the adjacent ring, an optionally substituted $\text{C}_5\text{--C}_6$ cycloalkyl or an optionally substituted $\text{C}_5\text{--C}_6$ aromatic ring;

R_5 and R_7 are independently H, or $\text{C}_1\text{--C}_8$ alkyl;


R_6 is selected from the group consisting of H, or $\text{C}_1\text{--C}_8$ alkyl, $\text{C}_2\text{--C}_8$ alkenyl and $\text{C}_2\text{--C}_8$ alkynyl;

15

n is an integer ranging from 0-2; and m is an integer ranging from 0-6.

In one embodiment a method of inhibiting angiogenesis comprises administering a compound having the general structure :


20




wherein R_1 is H, halo or $\text{C}_1\text{--C}_4$ haloalkyl;

R_2 is selected from the group consisting of

25



30



-16-

wherein R_{10} and R_{11} are independently selected from the group consisting of H, halo, C_1 - C_4 haloalkyl, $-NR_5R_7$, hydroxy and C_1 - C_8 alkoxy;

R_5 and R_7 are independently H, or C_1 - C_4 alkyl;

R_{12} is selected from the group consisting of H, halo and C_1 - C_8 alkoxy; and

5 R_6 is C_1 - C_8 alkyl. In one embodiment R_{10} and R_{11} are independently H, halo or C_1 - C_4 alkoxy, R_1 and R_{12} are H, and R_6 is C_1 - C_6 alkyl.

In accordance with one embodiment the thalidomide derivative compounds of the present invention can be formulated as pharmaceutical compositions by combining the compounds with one or more pharmaceutically acceptable carriers. These formulations can be administered by standard routes. In
10 general, the combinations may be administered by the topical, transdermal, oral, rectal or parenteral (e.g., intravenous, subcutaneous or intramuscular) route. In addition, the combinations may be incorporated into biodegradable polymers allowing for sustained release of the compound, the polymers being implanted in the vicinity of where drug
15 delivery is desired, for example, at the site of a tumor. Biodegradable polymers suitable for use with the present invention are known to the skilled practitioner and are described in detail, for example, in Brem et al., J. Neurosurg. 74:441-446 (1991).

In addition to the use of such pharmaceutical compositions as anti-angiogenic compounds, the thalidomide derived compounds and corresponding
20 compositions also have utility as sodium channel blockers, calcium channel blockers, contraceptives, anti-inflammatory agents and anti-cancer agents. In one embodiment a composition comprising a thalidomide derivative of the present invention is used to treat age related macular degeneration.

In accordance with one embodiment the present composition are
25 administered either orally or parenterally. When administered orally, the compounds are administered as a liquid solution, powder, tablet, capsule or lozenge. The compounds can be used in combination with one or more conventional pharmaceutical additives or excipients used in the preparation of tablets, capsules, lozenges and other orally administrable forms. When administered parenterally, and
30 more preferably by intravenous injection, the derivatives of the present invention can be admixed with saline solutions and/or conventional IV solutions.


-17-

The dosage of the active compound will depend on the condition being treated, the particular compound, and other clinical factors such as weight and condition of the human or animal and the route of administration of the compound. It is to be understood that the present invention has application for both human and veterinary use. For oral administration to humans, a dosage of between approximately 0.1 to 300 mg/kg/day, preferably between approximately 0.5 and 50 mg/kg/day, and most preferably between approximately 1 to 10 mg/kg/day, is generally sufficient.


It should be understood that in addition to the active anti-angiogenic compounds, the compositions of the present invention may include other agents conventional in the art including solubilizing agents, inert fillers, diluents, excipients and flavoring agents.

In accordance with one embodiment, diseases associated with corneal neovascularization can be treated by administering a composition comprising a compound having the general structure:

15



20




25


wherein R_1 is selected from the group consisting of H, halo, alkyl, haloalkyl, $-NR_5R_6$, hydroxy and alkoxy;

R_2 is selected from the group consisting of

-18-




5



R_3 is selected from the group consisting of

10



R_8 is selected from the group consisting of H and alkyl, or R_3 and R_8 taken together, can form with the adjacent ring, an optionally substituted C_5 - C_6 cycloalkyl or C_5 - C_6 aromatic ring;

R_{10} and R_{11} are independently selected from the group consisting of H, halo, C_1 - C_4 haloalkyl, $-NR_5R_7$, hydroxy and C_1 - C_8 alkoxy or R_{10} and R_{11} taken together, can form with the adjacent ring, an optionally substituted C_5 - C_6 cycloalkyl or optionally substituted C_5 - C_6 aromatic ring;

20


R_5 and R_7 are independently H, or C_1 - C_4 alkyl;

R_{12} is selected from the group consisting of H, halo and C_1 - C_8 alkoxy;

m is an integer ranging from 0-4; and

R_6 is C_1 - C_8 alkyl. In one embodiment the compound has the general structure

25




wherein R_1 is H or halo; and

30

R_2 is selected from the group consisting of

-19-



wherein R_{10} and R_{11} are independently H, halo or C_1 - C_4 alkoxy, R_1 and R_{12} are H, and R_6 is C_1 - C_6 alkyl.

Another disease which can be treated according to the present invention is rheumatoid arthritis. It is believed that the blood vessels in the synovial lining of the joints undergo angiogenesis. In addition to forming new vascular networks, the endothelial cells release factors and reactive oxygen species that lead to pannus growth and cartilage destruction. The factors involved in angiogenesis may actively contribute to, and help maintain, the chronically inflamed state of rheumatoid arthritis.


The thalidomide derived compounds of the present invention are also anticipated to have use in treating a wide variety of diseases or conditions that are related to angiogenesis, including diabetic retinopathy, retinopathy of prematurity, corneal graft rejection, neovascular glaucoma and retrolental fibroplasia, epidemic keratoconjunctivitis, Vitamin A deficiency, contact lens overwear, atopic keratitis, superior limbic keratitis, pterygium keratitis sicca, sjogrens, acne rosacea, phlyctenulosis, syphilis, Mycobacteria infections, lipid degeneration, chemical burns, bacterial ulcers, fungal ulcers, Herpes simplex infections, Herpes zoster infections, protozoan infections, Kaposi sarcoma, Mooren ulcer, Terrien's marginal degeneration, mariginal keratolysis, trauma, rheumatoid arthritis, systemic lupus, polyarteritis, Wegeners sarcoidosis, Scleritis, Steven's Johnson disease, pemphigold radial keratotomy, and corneal graph rejection. Diseases associated with retinal/choroidal neovascularization that can also be treated according to the present invention and include, but are not limited to, diabetic retinopathy, macular degeneration, sickle cell anemia, sarcoid, syphilis, pseudoxanthoma elasticum, Pagets disease, vein occlusion,

-20-

artery occlusion, carotid obstructive disease, chronic uveitis/vitritis, mycobacterial infections, Lyme's disease, systemic lupus erythematosus, retinopathy of prematurity, Eales disease, Bechets disease, infections causing a retinitis or choroiditis, presumed ocular histoplasmosis, Bests disease, myopia, optic pits, Stargarts disease, pars
5 planitis, chronic retinal detachment, hyperviscosity syndromes, toxoplasmosis, trauma and post-laser complications.


Example 1

Molecular modeling was used to facilitate the design of a novel series
10 of isoquinolines. Initially, a one carbon addition to the imide ring was considered to prepare compounds of the general structure:



15 Conformational analysis of this phthalimide analog, docked in the crystal structure of polyADP-ribose polymerase, revealed the conformational similarity of the proposed structural mimic with phthalimides. With this in mind, the synthesis of two initial series of analogs was proposed as a preliminary set of synthetic targets.

20




10

compound	R ₂₀	R ₂₁	amino acid	n
Series A				
1	H	H	glycine	
2-(±)	CH ₃	H	alanine	
3-(±)	benzyl	H	phenylalanine	
4-(±)	HO-benzyl	H	tyrosine	
5	phenyl	phenyl		
6-(±)	phenyl	H		
7-(±)	phenyl	CH ₃		
Series B				
8				2
9				3
10				4

15


20

These compounds represent first generation analogs which investigate important structure and enantioselective relationships for isoquinolines. The synthesis of mimics **1-10** is outlined in Scheme I and represents reasonable literature reported organic transformations. The utility of this synthetic scheme allows for the insertion of a large diversity of functional groups. Furthermore the synthetic design allows for the straightforward synthesis of enantiomerically pure final products through the addition of selected chiral amino acids.

Scheme I. Synthetic Scheme for Analogs in this Study**Example 2**

Additional compounds were prepared wherein the piperidine-2,6-dione moiety of thalidomide is replaced with a rigid or open ring structure having the general structure:


15




20 wherein R_{22} is aryl and R_{23} is H, C_1 - C_{12} alkyl, C_2 - C_{12} alkenyl or C_2 - C_{12} alkynyl, or R_{22} and R_{23} together with the intervening carbons form an aryl. The synthetic schemes used to prepare these compounds is as follows:

-23-


Synthesis of 2-Indan-5-yl-1,3-dioxo-2,3-dihydro-1H-isoindole-5-carboxylic acid




15 Compounds 5-7 can also be prepared by the following reaction:



Synthesis of 2-(1,2,3,4-Tetrahydro-naphthalen-1-yl)-isoindole-1,3-dione



-24-



12-15


16-22

Compd #	R ₂₄	n	yield	Compd#	R ₂₄	R ₂₅	n	yield
12	H	1		16			1	98%
13	H	3	70%	17	H	H	3	100%
14	H	3		18	H	3-CH ₃	3	76%
15	3-F	3	48%	19	3-F	2-F	3	62%
				20	3-F	3-CH ₃	3	50%
				21	H	2-F	3	72%
				22	H	H	2	84%

15


Synthesis of

2-(10,11-Dihydro-5H-dibenzo[a,d]cyclohepten-5-yl)-isoindole-1,3-dione




25

30




-25-


Synthesis of 2-(1-Naphthalen-2-yl-propyl)-isoindole-1,3-dione





Synthesis of 2-[1-(3,4-Dimethoxy-phenyl)-heptyl]-isoindole-1,3-dione



-26-

Synthesis of 2-Isoquinolin-1-yl-isoindole-1,3-dione



10 **Synthesis of N-(2-phenylethyl)-phthalimide**

**Synthesis of Phenyl Substituted-phthalimides**

30


-27-

Synthesis of 4-Hydroxy-2H-isoquinolin-1-one



15


Synthesis of Pyrrolo[1,2-b]isoquinoline-5,10-dione



25

-28-

Synthesis of 3H-quinazolin-4-ones



These compounds will be evaluated for their ability to inhibit corneal neovascularization and endothelial cell proliferation.

5

Example 3

Endothelial cell proliferation assay.











The human vascular endothelial cells (HUVECS) were cultured to peri-confluence (80%) in 20% serum and treated with thalidomide (standard) or its analog (40-400 μ M). After 20 h, [³H]-thymidine (2 μ Ci/ml) was added to the culture medium for 2-4 h. The [³H]-thymidine incorporation was stopped with ice-cold PBS (3 washes) and the cells were incubated with cold 10% trichloroacetic acid (TCA) for 10 min at 4 °C. The cells were further incubated with TCA at room temperature for 10 min and washed three times with PBS. The cells were solubilized overnight with 1N NaOH and neutralized with an equivalent amount of 1N HCl before radioactivity was determined. The anti-proliferative activity of thalidomide or its analog was computed as a percent inhibition of HUVECS mitogenic response to 20% serum (fetal calf serum).

Evaluation of thalidomide for inhibition of the incorporation of ³H-thymidine into endothelial cells reveal this compound to have an IC₅₀ of 200 μ M. Preliminary IC₅₀ data for the first panel of compounds reveal four compounds (5, 29, 32 and 50) to have 2-4 times greater potency than thalidomide in inhibiting ³H-thymidine incorporation into endothelial cells (See Table 1). This structure activity

20

relationship (SAR) establishes that the piperidine-2,6-dione moiety is not needed for increased inhibition of endothelial cell proliferation.

Table 1. Biological Data for Analogs in this Study


5	Compd #	Structure	Inhibition of Endothelial Cell Proliferation (IC ₅₀ μM)
	1	 Thalidomide	200
10	5		85
	7		>400
	49		385
15	29		45
	32		<38.5
20	50*		110
	51*		350
	52*		>400
25	53*		>400

*Obtained from the Aldrich Chemical Library.


-30-

Example 4

4-oxo-quinazoline derivatives having the general structure:





wherein n is 1-4 can be prepared according to the following schemes:





Compound	chiral center	R ₃₀	R ₃₁	% yield
33		H	H	76
34	R	C ₆ H ₅	H	83
35	S	C ₆ H ₅	H	75
36	S	HOC ₆ H ₄	H	--
37	R	CH ₃	H	42
38	S	CH ₃	H	68
39	S	CH ₂ C ₆ H ₄	H	90
40	R	CH ₂ C ₆ H ₄	H	16

-31-




15 Preparation of expanded ring structures:




Claims

1. A compound represented by a general structure selected from the group consisting of:




wherein R_1 is selected from the group consisting of H, halo, C_1 - C_4 alkyl, carboxy and C_1 - C_8 alkoxy;

R_2 is selected from the group consisting of



R_3 is selected from the group consisting of



R_8 is selected from the group consisting of H, C_1 - C_8 alkyl, phenyl and benzyl or R_3 and R_8 taken together, can form, with the adjacent ring, an optionally substituted 5- or 6-membered heteroaryl;

30 R_9 and R_{13} are independently selected from the group consisting of halo, methoxy and C_3 - C_8 alkoxy;

R_{10} and R_{11} are independently selected from the group consisting of H, halo,

-33-

C₁-C₈ alkyl, C₁-C₈ haloalkyl, -NR₅R₇, hydroxy and C₁-C₈ alkoxy or R₁₀ and R₁₁ taken together, can form with the adjacent ring, an optionally substituted C₅-C₆ cycloalkyl or an optionally substituted C₅-C₆ aromatic ring;

R₅ and R₇ are independently H, or C₁-C₈ alkyl;

5 R₆ is selected from the group consisting of H, or C₁-C₁₂ alkyl, C₂-C₈ alkenyl and C₂-C₈ alkynyl;

R₁₂ is selected from the group consisting of H, halo, C₁-C₄ haloalkyl, -NR₅R₆, C₁-C₈ alkyl, hydroxy and C₁-C₈ alkoxy;


n is an integer from 1-3 and m is an integer ranging from 0-6.

10


2. The compound of claim 1 wherein

R₁ is H or halo;

R₂ is selected from the group consisting of



R₃ is selected from the group consisting of



R₄ and R₈ are H;

R₆ is C₁-C₈ alkyl;

25 R₁₀ and R₁₁ are independently selected from the group consisting of H, halo, and C₁-C₈ alkoxy or R₁₀ and R₁₁ taken together, can form with the adjacent ring, an optionally substituted C₅-C₆ cycloalkyl or an optionally substituted C₅-C₆ aromatic ring;


R₁₂ is selected from the group consisting of H, halo, and C₁-C₄ alkoxy; and

30

m is 1 or 0.


-34-

3. The compound of claim 1 wherein the compound is represented by the general structure:



wherein R_1 is H or halo; and

R_2 is selected from the group consisting of




wherein R_6 is H, or C_1 - C_8 alkyl; and

R_{10} and R_{11} are independently selected from the group consisting of H, halo, $-NH_2$, hydroxy and C_1 - C_8 alkoxy.

4. The compound of claim 3 wherein R_1 is H.


5. A pharmaceutical composition comprising the compound of claim 1 and a pharmaceutically acceptable carrier.

6. A method of inhibiting angiogenesis in a warm blooded vertebrate, said method comprising the steps of administering to said vertebrate a composition comprising a compound of the general structure




-35-

wherein R_2 is selected from the group consisting of



R_3 is selected from the group consisting of



R_8 is selected from the group consisting of H, C_1 - C_8 alkyl, phenyl, and benzyl or R_3 and R_8 taken together, can form, with the adjacent ring, an optionally substituted 5- or 6-membered heteroaryl;

20 R_{10} and R_{11} are independently selected from the group consisting of H, halo, C_1 - C_8 alkyl, C_1 - C_8 haloalkyl, $-\text{NR}_5\text{R}_7$, hydroxy and C_1 - C_8 alkoxy or R_{10} and R_{11} taken together, can form with the adjacent ring, an optionally substituted C_5 - C_6 cycloalkyl or an optionally substituted C_5 - C_6 aromatic ring;

R_5 and R_7 are independently H, or C_1 - C_8 alkyl;

25 R_6 is selected from the group consisting of H, or C_1 - C_{12} alkyl, C_2 - C_8 alkenyl and C_2 - C_8 alkynyl;

R_{12} is selected from the group consisting of H, halo, C_1 - C_4 haloalkyl, $-\text{NR}_5\text{R}_6$, C_1 - C_8 alkyl, hydroxy and C_1 - C_8 alkoxy; and


m is an integer ranging from 0-6.

30

-36-

7. The method of claim 6 wherein the compound has the general structure:


5



wherein R_1 is H or halo; and

R_2 is selected from the group consisting of

10



wherein R_6 is H, or C_1 - C_8 alkyl; and


15

R_{10} and R_{11} are independently selected from the group consisting of H, halo, and C_1 - C_8 alkoxy or R_{10} and R_{11} taken together, can form with the adjacent ring, an optionally substituted C_5 - C_6 cycloalkyl or an optionally substituted C_5 - C_6 aromatic ring.


20

8. A pharmaceutical composition comprising a compound represented by the general structure:

25





30




wherein R_1 is H or halo;

-37-

R_2 is selected from the group consisting of



R_3 is selected from the group consisting of



R_{10} and R_{11} are independently selected from the group consisting of H, halo, $\text{C}_1\text{--C}_8$ alkyl, $\text{C}_1\text{--C}_8$ haloalkyl, $\text{---NR}_5\text{R}_7$, hydroxy and $\text{C}_1\text{--C}_8$ alkoxy or R_{10} and R_{11} taken together, can form with the adjacent ring, an optionally substituted $\text{C}_5\text{--C}_6$ cycloalkyl or an optionally substituted $\text{C}_5\text{--C}_6$ aromatic ring;


R_5 and R_7 are independently H, or $\text{C}_1\text{--C}_8$ alkyl;

R_6 is selected from the group consisting of H, or $\text{C}_1\text{--C}_{12}$ alkyl, $\text{C}_2\text{--C}_8$ alkenyl and $\text{C}_2\text{--C}_8$ alkynyl;

R_{12} is selected from the group consisting of H, halo, $\text{C}_1\text{--C}_4$ haloalkyl, $\text{---NR}_3\text{R}_6$, $\text{C}_1\text{--C}_8$ alkyl, hydroxy and $\text{C}_1\text{--C}_8$ alkoxy; and

m is an integer ranging from 0-2; and a pharmaceutically acceptable carrier.


8. A pharmaceutical composition comprising a compound represented by the general structure:



-38-

wherein R_1 is H or halo; and

R_2 is selected from the group consisting of



wherein R_6 is C_1 - C_8 alkyl; and

R_{10} and R_{11} are independently selected from the group consisting of H, halo, and C_1 - C_8 alkoxy.

- 15 9. The composition of claim 8 wherein R_1 is H and R_{10} and R_{11} are independently selected from the group consisting of H and C_1 - C_8 alkoxy.

20